

# Common themes in geriatric emergency general surgery: a narrative review

Giorgia Santandrea,<sup>1</sup> Carlo Vallicelli,<sup>1</sup> Paolo Carcoforo,<sup>2</sup> Silvia Barbara,<sup>2</sup> Vanni Agnoletti,<sup>3</sup> Francesca Bravi,<sup>4</sup> Fausto Catena<sup>1</sup>

<sup>1</sup>Emergency and Trauma Surgery Department, Bufalini Hospital, Cesena; <sup>2</sup>Department of Morphology, Experimental Medicine and Surgery, University of Ferrara; <sup>3</sup>Anesthesia, Intensive Care and Trauma Department, Bufalini Hospital, Cesena; <sup>4</sup>Healthcare Administration, Santa Maria delle Croci Hospital, Ravenna, Italy

## Abstract

Population aging is a growing phenomenon. The geriatric patient is defined by physical, psychological, and social characteristics associated with aging that make him or her more fragile and susceptible. An older patient is usually fragile: frailty has a linear, if not unambiguous, relationship with age. With this background, the concept of futility emerges in emergency surgery, which refers to having a goal but being almost certain of failure in achieving it. A narrative review of the literature was carried out using PubMed, Google Scholar, and Cochrane to identify pertinent publications. Inclusion criteria included: i) an emergency setting in geriatric general surgery, ii) frailty in emergency surgery, and iii) futility in emergency surgery. The study identified valid assessment scores for older and frail patients, as well as imaging tools that may aid in the

evaluation of frailty, demonstrating the potential futility of surgical treatment. The review addressed the most common acute surgical situations that affect geriatric patients. Older patients are more fragile and vulnerable than the general population, even though geriatric age does not always correlate with frailty. Several fragility scores have been developed to perform an accurate preoperative assessment of the urgent elderly surgical patient, to estimate treatment futility and perioperative risk, and to guide the surgeon to the most appropriate therapeutic or palliative action.

## Introduction

Population aging is a constantly increasing phenomenon. The geriatric patient (conventionally defined as a person over 65 years of age) is characterized by physical, psychological, and social traits that make him or her more fragile and vulnerable compared to the general population. Advanced age is associated with an increased risk of comorbidities and loss of autonomy, as well as a reduced life-expectancy. Aging refers to the inevitable and irreversible organ malfunction with time, regardless of the absence of acute organ damage, disease, or unbalanced lifestyles. Typical signs of aging can be summarized as a reduced ability to maintain homeostasis in response to stressful events and therefore a limited proper reaction to acute events, with subsequent increased risks of morbidity, loss of independence during ADLs (Activities of Daily Living), and mortality. The concept of “geriatric age”, however, must take into account not only the patient’s chronological age, but also social, economic, and cultural factors, including the patient’s functional status. For this reason, it would be more appropriate, in the overall assessment of a patient, to speak of “frailty” instead of geriatric age.

Frailty is defined as a reduced physiological reserve associated with an increased vulnerability due to the gradual loss of physical, cognitive, social, and psychological functions, that leads to a higher susceptibility to adverse events, disability, and death.<sup>1</sup> Frailty displays a linear, though not univocal, trend with age. In surgery, frailty is related to an increased risk of all complications (clinical and surgical) and death: the patient’s diminished resources cause an inadequate ability to promptly respond to surgery, and the presence of comorbidities raises the risk of complications and death in the post-operative period, as well as dampen the patient’s abilities and autonomy. Even though in elective surgery it’s feasible to perform a multidimensional assessment of the geriatric patient and adequately prepare him to face the surgical stress through pre-operative-habilitation programs, this is very complicated and often not possible in urgent or emergent situations.

As a matter of fact, in an emergent surgery setting, the patient’s assessment and treatment choice must be as hasty as possible, but the critically ill patient might have a physical or sensory impairment, and his/her family members may not be immediately avail-

Correspondence: Giorgia Santandrea, Emergency and Trauma Surgery Department, Bufalini Hospital, 47521 Cesena, Italy.  
E-mail: giorgia.santandrea@auslromagna.it

Key words: futility; geriatric patient; emergency surgery.

Conflict of interest: the authors declare no potential conflict of interest, and all authors confirm accuracy.

Ethics approval: not applicable.

Informed consent and patient consent for publication: not applicable.

Availability of data and materials: all data generated or analyzed during this study are included in this published article.

Received: 7 January 2024.

Accepted: 21 May 2024.

Publisher’s note: all claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

©Copyright: the Author(s), 2024

Licensee PAGEPress, Italy

Surgery in Geriatrics and Frailty 2024; 1:11

doi: 10.4081/sigaf.2024.11

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial International License (CC BY-NC 4.0) which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

able; this may make it difficult to carry out an overall assessment of the patient and the risks connected with a possible surgical treatment. At least 50 different scores have been suggested and are currently being used in surgical emergencies in frail patients, but an adequate pre-operative assessment in these cases is often very complex.

Thus the concept of futility in emergency surgery. Futility is the lack of effectiveness; it consists of setting a goal and carrying out the action(s) aimed at achieving it, but with the awareness that the aforementioned action(s) will be useless and unsuccessful.

This notion is corroborated by the acknowledgment of the inherent risk in a clinical case where uncertainty and the danger of morbidity and death are extreme. The evaluation of futility in emergency surgery unfolds several considerations, from the strictly clinical ones (e.g. the overall risks for the patient in question) to the psychological and social ones (e.g. the loss of autonomy in daily activities, or the worsening in the quality of life), and to the ethical ones (e.g. the decision not to proceed surgically as it would be 'futile'). It is in such a complex and multifactorial background that it was mandatory to examine several different variables and develop the tools to shape a patient's global assessment as objectively as possible.

This literature review aims to analyze and discuss what might be the best approach for the emergency and urgent surgeon in assessing the geriatric and fragile patient and therefore in deciding whether or not a particular surgical act is useful.

## Materials and Methods

Literature research was performed using PubMed, Google Scholar, and Cochrane to identify relevant publications, using "emergency" AND "general surgery" AND "geriatric" as search terms. The search was limited to publications written in English, rejecting duplicates and selecting a total of 98 publications (Figure 1). Abstracts were screened for inclusivity and full articles were reviewed to identify all potentially relevant studies. The inclusion criteria were: i) emergency in geriatric general surgery, ii) frailty in emergency surgery, and iii) futility in emergency surgery. Studies dealing with frailty in disciplines other than general surgery and with emergency surgery in the non-geriatric patient setting were excluded.

It was not possible to develop a systematic literature review with meta-analysis because of the heterogeneity of the studies, the outcomes currently under review, and the ample variability of approach to the different outcomes in the analyzed publications.




Figure 1. Publications selection.

## Futility

The concept of futility consists of setting a goal and carrying out the actions and activities aimed at achieving that goal, but with the virtual assurance that the aforementioned actions will fail to achieve it.<sup>3,4</sup> Futility is associated with an intrinsic risk in an event where the danger and uncertainty for the patient's outcomes are extreme. Although risk evaluation is vital even in elective surgery settings, it becomes crucial in emergency contexts as these are associated with an increased complexity of the patient's global assessment related to the need for swift decisions, often made late at night, a narrower window for adequate multidisciplinary discussion, and possible lack of family support to guide the surgical decision.

The concept of futility can be described as i) quantitative: *i.e.*, as the statistical probability of surviving a specific procedure, or as the probability of success of a treatment;<sup>5,6</sup> ii) qualitative: *i.e.*, as the benefit that the procedure may bring regarding the patient's quality of life (e.g. the presence of a permanent ostomy, or the need for palliative therapy in oncology).<sup>5,6</sup>

The analysis of both these traits can guide the discussion and risk stratification in clinical decision-making. Several parameters help predict the futility of a surgical procedure: advanced age, serum lactates, renal function, patient's comorbidities, state of consciousness, and the presence or absence of septic status. The quantitative feature of futility is the one that predominantly leads the clinical decision, however, it may not always be easily understood or acknowledged by the patient or his/her family, as the qualitative aspects related to the procedure may often be more meaningful for the patient and his/her family than the risk of death itself. To exemplify, for patients it's easier to understand the risk of being discharged with a permanent ostomy or the need for home support than the risk of death related to their clinical condition.<sup>7</sup>

These considerations clarify the extreme controversy of the term futility in the decision-making process and the need for an equilibrium between the urgent nature of the surgical act and its associated risk, as emergency surgery increases the risk of surgery without gross benefits (*i.e.*, death within 48-72 hours after surgery).

The overall management of these complex scenarios can turn out to be pretty difficult for the surgeon, thus leading to emotional stress, depression, and burnout.<sup>8</sup> The first step should be to balance the surgical risk using objective scores combined with the patient's values and perception. The complementary assessment of the risks, benefits, and therapeutic alternatives can then guide the conversation to a decision. A multidimensional assessment between the anesthetist, the surgeon, and the patient's physician, if possible, could be valuable, as it would provide the patient and his family a wide overview of potential surgical benefits, associated risks of morbidity and mortality, potential loss of autonomy in daily activities, and possible non-operative alternatives.<sup>9</sup>

## Geriatric patient evaluation

Geriatric age conventionally begins at 65 years of age, but some patients may require earlier geriatric expertise because of underlying conditions and comorbidities that make them frail. Aging refers to the inevitable and irreversible organ decline with the passing of years, even in the absence of acute damage, diseases, environmental hazards, or unbalanced lifestyles. The first sign of aging corresponds to a reduced ability to maintain homeostasis under stress, underlying thus frailty in the elderly. Frailty indeed shows a linear, though not univocal, trend with age: it is reckoned that frailty is present in 10-

20% of patients older than 65 years of age, and this percentage increases up to 40% in patients older than 80 years of age.<sup>10</sup> Disease interferes with natural aging and triggers specific geriatric complications, known as geriatric syndromes (*i.e.*, delirium, incontinence, falls, visual and hearing impairments, and weight loss).<sup>11</sup>

Emergency general surgery in older patients is associated with increased mortality, and higher rates of short- and long-term complications (*i.e.*, in emergency cholecystectomy, complications can reach up to 27% in 65-79 year-old patients, and up to 38% in patients aged 80 years or older),<sup>12</sup> loss of independence and autonomy. The need to compute risk stratification, throughout common scores, also applied in elective surgery (ASA, APACHE II, P-POSSUM, etc),<sup>13</sup> frailty assessment, and presence/absence of sarcopenia are hence vital.

### ASA-score (American Society of Anesthesiologists Physical Classification System)

This scale is effective in establishing how fit the patient is for a surgical procedure. It classifies comorbidities, but lacks information on operative risk, as it should be merged with information about frailty, type of surgery, and available medical facilities. Moreover, it does not consider age and physical status, as well as other comorbidities such as cancer, and may be biased towards the clinician's subjective judgment. In an emergency setting in a geriatric patient, the score appears adequate for predicting mortality, but less suitable for estimating short-term complications.<sup>14-17</sup>

### APACHE II scoring system (Acute Physiology and Chronic Health Assessment)

It is a measuring system based on current physiologic measurements, age, and previous health conditions, often used in the ICU setting, determines disease severity in critically ill patients within the first 24 hours of intensive care hospitalization. In an emergency surgery setting in the elderly, this score is both inadequate in predicting short-term and long-term complications as well as time-consuming and onerous to perform.<sup>12,14,17,18</sup>

### CFS (Clinical Frailty Scale)

It is a simple and quickly validated scale to assess patients' frailty and fitness, tailored for patients aged 65 years or older, administered on triage admission and then 2 weeks later. It labels the population into 9 groups based on comorbidities, physical activity, and autonomy in activities of daily living. It's reliable in predicting short- and long-term mortality, but not in estimating short-term surgical complications.<sup>14,17,19-25</sup>

### ACS-NSQIP risk calculator (American College of Surgeons National Surgical Quality Improvement Program)

This is an online calculator that takes into account the patient's age, comorbidities (obesity, diabetes, cardiovascular disease, dyspnea, hepatopathy, renal failure, advanced neoplasm), medication use (*i.e.*, corticosteroids or antihypertensive drugs), the patient's lifestyle (smoker, drinker), and the type of surgery planned. These data are enriched with information about the geriatric patient's independence status. It measures the risk of complications within the first 30 days after surgery and thus allows an accurate prediction of short-term complications and death risks; it may be useful guidance in the surgical-decision making.<sup>14</sup>

### P-POSSUM score (Porthsmouth Physiological Operative Severity Score for enUmeration of Mortality and morbidity)

It is a verified score that allows a projection of morbidity and mortality risks in the surgical patient based on objective physiological and operative criteria; it is therefore a great assessment tool for the frail patient even in the emergency setting.<sup>12,17,26</sup>

### NELA Risk calculator

It provides an estimate of the risk of death within 30 days of emergency abdominal surgery. It has been developed using the data gathered from patients admitted into the National Emergency Laparotomy Audit between December 2016 and November 2019. This tool reviews each procedure's risks and patient's information. In the UK it has proved its superior accuracy over the P-POSSUM score for patients undergoing emergency laparotomy.<sup>36</sup>

### Comprehensive geriatric assessment

Preoperative patient stratification may be difficult in an emergency setting; risk assessment in older patients in emergency surgery is however cardinal for a more appropriate intraoperative physiological control and optimal post-operative management. The concept of Comprehensive Geriatric Assessment (CGA) is increasingly emphasized in current literature as the standard key for improving outcomes in elderly patients in elective surgery.<sup>34-37</sup> The CGA consists of a global evaluation of the geriatric patient, taking into account age, comorbidities, degree of autonomy in ADLs, psychological cognitive and social status, and family support, thus allowing not only to rate his or her surgical risk, but also to organize postoperative recovery through a targeted physiotherapy program, to adequately plan hospital discharge (towards home or care facilities), and, in case of home discharge, to guide family members in the overall assistance and management of any presides. This assessment should be managed by a multidisciplinary team (composed usually of a geriatrician, a nurse, and a social worker) so that all areas of concern can be suitably and accurately determined. Other figures, such as a psychiatrist or psychologist, may be involved. Predefined questionnaires are employed to inquire about the ability to perform functional actions and the possible need for assistance, history of falls, presence of urinary and/or fecal incontinence, pain, available social support, depressive symptoms, vision or hearing impairment, nutritional status, comorbidities, and polypharmacotherapy. Following the analysis of these parameters, the multidisciplinary team then defines the most appropriate setting for the patient (home, a long-term care facility, a hospital ward) and any necessary supplies. In a surgical urgency/emergency setting, a preoperative multidisciplinary evaluation of the patient is impractical due to the acute nature of the situation; it would be desirable however to perform it routinely both in the post-operative period and in case of conservative palliative treatment.

### Frailty evaluation

Frailty is defined as a condition of reduced physiological reserve associated with enhanced vulnerability, due to a loss of physical, cognitive, social, and psychological functioning, leading to a raised susceptibility to adverse events, disability, and death.<sup>1,2</sup> Frailty, in the surgical setting, is related to an increased rate of all

complications and death:<sup>37</sup> frail patients' limited resources result in a decreased ability to respond adequately to surgery; moreover, potential comorbidities are linked with a higher risk of postoperative complications.<sup>12,39,40</sup> The frail surgical patient has indeed, compared with the non-frail patient, an increased risk of developing complications, both surgical and medical (e.g., wound infections, reintervention, pneumonia, renal failure), as well as an amplified postoperative disability and mortality,<sup>11,19,34,41</sup> longer length of hospital stay, increased institutionalization, and higher rates of hospital readmission.

Frailty is not an absolute contraindication to surgery, but it is related to an increased intra and postoperative risk;<sup>2,42</sup> risk stratification can therefore be convenient in predicting outcomes and optimizing patients when faced with surgical stress while trying to avoid the patient's functional decline, prevent or manage appropriately early postoperative complications, plan the post-surgical course and proper setting after hospital discharge.<sup>39,40</sup> Plentiful scores have been developed to optimize the frail patient's assessment in an urgent/emergent surgical setting. There are around fifty such scores, a few examples of those most commonly used in an emergency/emergency setting are the CFS (Clinical Frailty Score),<sup>12,19-25</sup> which we have already discussed before, and the following ones.

### Fried's Frailty Index

This score was issued from cardiovascular health-related studies. It examines 5 parameters: unintentional weight loss, gait slowness, easy fatigability, history of falls, and reduced strength in grasping objects. The patient is considered a frail individual when the score is greater than or equal to 3. The advantage of this score in an emergency setting appears limited, as the patient's dire condition may override the assessment.<sup>12,44,45</sup>

### mFFC (modified Fried's Frailty Criteria)

This is a multidimensional frailty screening system, similar to the Fried Index, from which it derived, that assesses grip strength, fatigability, scant physical activity, weight loss, and exhaustion. A score of 0 identifies a robust patient, a score between 1-2 a pre-frail patient, and a score greater than or equal to 3 identifies a frail patient. Again, the worth of this index may once more be distorted in an emergency.<sup>36</sup>

### mFI-11 (modified Frailty Index 11)

This score takes into account the patient's comorbidities for a total of 11 points: i) diabetes mellitus, ii) congestive cardiac failure, iii) medically treated hypertension, iv) history of stroke or TIA, v) partial or total dependence in ADLs, vi) history of myocardial infarction, vii) history of peripheral vascular disease, viii) history of cerebrovascular pathology with relics, ix) COPD or pneumonia, x) history of PTCA or angina, and xi) history of sensory impairments. A point is assigned to each variable: if the score is 0 the patient is considered non-fragile, if the score is 1-2 the patient is considered pre-fragile, and above 3 points the patient is considered frail. A higher score is associated with increased complications, such as wound infections, reintervention, hospital readmission after discharge, increased institutionalization, longer hospital stay, and worse preoperative mortality.<sup>2,41</sup>

### mFI-5 (modified Frailty Index-5)

This modified score is more flexible and brief than the mFI-11, and is obtained with only 5 elements: i) presence of heart failure

within 30 days of surgery, ii) diabetes mellitus, iii) COPD or pneumonia, iv) partial or total dependence in ADLs, v) treated hypertension. A score of 1-2 suggests intermediate frailty, which has a 1.5% increased risk of failure-to-rescue, postoperative complications, reintervention, and mortality; a score greater than or equal to 3 reflects severe frailty, resulting in a 4-fold increase in all-cause mortality, a 5-fold increase in perioperative complications, and a 2-fold increase in reintervention risk.<sup>41,42</sup>

### fTRST (flamish version of the Triage Risk Screening Tool)

This is a 5-item questionnaire, with a total score ranging from 0 to 6, which allows for an overall assessment of the patient. It considers cognitive decline, autonomy in ADLs, mobility and any falls in the past 6 months, history of hospitalization in the past 3 months, and poly-pharmacotherapy (more than 5 chronic medications). A score greater than or equal to 2 correlates with enhanced short- and long-term postoperative complications, higher mortality, and longer duration of hospital stay. It's a rather simple and quick tool;<sup>19,20,40,46</sup>

### 50-RPFI (modified 50-variable Rockwood Preadmission Frailty Index)

It consists in a 50-item score that examines comorbidities, ADLs and IADLs, psychological, nutritional, and functional status, and a few blood test values. It's very useful in assessing the risk of postoperative morbidity and mortality, the main drawback is that it is quite elaborate and long to complete;<sup>20</sup>

### TSFI (Trauma Specific Frailty Index), EGSFI (Emergency General Surgery Frailty Index), and TEGS-FI (Trauma and Emergency General Surgery Frailty Index)

TSFI<sup>1,47</sup> is the only model developed specifically to assess frailty in the trauma patient; its application was later extended to emergency surgery (EGSFI),<sup>1,20,48,49</sup> hence TEGS-FI. The score is characterized by 15 variables and examines the presence of comorbidities (neoplasm, cardiovascular disease, dementia), attitude towards health status, ADLs, and functional and nutritional status. A higher score is linked to an increased risk of complications, but its regular employment allows an overall reduction in the number of complications and institutionalization, whilst it doesn't affect mortality.<sup>50</sup>

### MALE risk score

This score weighs up factors such as male sex (male), presence of anemia (anemic), albuminemia  $\leq 3.5$  g/dl (low albumin), age greater than or equal to 85 years (Eighty-five years old); each variable is assigned one point. This is a practical, easy-to-calculate, and accurate tool for predicting poor outcomes in elderly patients in an emergent surgery setting. Indeed, in a prospective multicenter study, Ablett *et al.*<sup>51</sup> imply that a higher MALE score is related to an increased 30- and 90-day mortality, new hospitalizations in patients who sustained emergent surgery, and longer length of hospital stay. The importance of hypoalbuminemia is especially emphasized: albumin values below 3.5 g/dl are associated with malnutrition states that may underlie chronic inflammatory bowel diseases, hepatopathy, neoplasms, and heart failure, thus associating with a higher rate of surgical complications, such as anastomotic dehiscences and wound infections.<sup>51,52</sup>

## Frailty imaging

Communication with the patient and/or family members in an emergency setting may be complicated, making it challenging to apply the available scores. CT-guided radiological methods have therefore been implemented to assess patients' characteristics suggestive of a potential vulnerability to the surgical act: i) sarcopenia: a progressive and generalized skeletal muscle syndrome characterized by loss of muscle mass, reduced strength, and muscle quality; it's calculated by measuring the TPI (Total Psoas Index), comprised of the sum of the area of the right and left psoas muscle at L3 level on CT, divided by the patient's height in meters;<sup>12,28,29</sup> ii) osteopenia: a measurement of attenuation, in Hounsfield Units (HU), in the region of interest (ROI). A 2D ROI is assessed at the level of the anterior vertebral trabecular area on an axial projection of L3. It has proved its usefulness in predicting 30-day mortality in patients requiring emergency laparotomy;<sup>28,30</sup> iii) sarcopenic obesity: the total visceral and subcutaneous adipose tissue area measured in an axial projection and divided by total muscle area at L3 level. There are limited data regarding the advantage of this parameter in assessing emergency morbidity and mortality, the main field of application concerns elective gastric and pancreatic surgery;<sup>27,31-34</sup> iv) renal volume normalized to the patient's height; v) calcification rate of the abdominal aorta: the aorta is segmented from D12 to the lower end of L3; the calcification rate is obtained by dividing the percentage of calcification volume by the total volume of the aorta.

The adequacy of these last two parameters in assessing frailty in older patients has been hypothesized, but there are only a handful of studies available.<sup>28</sup>

## Main emergency surgical diseases in older patients

### Acute cholecystitis

The MICOL study<sup>53</sup> shows that age is a strong risk factor for biliary cholecystitis in both sexes: the prevalence of cholelithiasis at 70 years of age is 15% in men and 24% in women, while in nonagenarians the prevalence reaches 24% in men and 35% in women. Clinical manifestations of acute cholecystitis in geriatric patients may be altered by factors such as reduced pain perception,<sup>54,55</sup> altered biliary physiology,<sup>56</sup> and a different response to tissue damage.<sup>57</sup> The typical symptom of acute cholecystitis, characterized by epigastric and right hypochondrium pain, may be absent in older patients, either as an effect of aging or as a consequence of pathologies such as diabetes, known precursor to an altered nociceptive sensitivity.<sup>13</sup> 12% of patients with acute cholecystitis experience atypical pain, while 5% of elderly patients are pain-free,<sup>58</sup> with an often negative Murphy's sign. Fever is reported with inconsistent percentages within the geriatric population (36-74%), but only 6-10% reach a body temperature above 38°C.<sup>58,59</sup> Regarding laboratory tests, only 41-59% of elderly patients with acute cholecystitis present with neutrophilic leucocytosis.<sup>58,60</sup> The diagnosis of acute cholecystitis may therefore often be delayed, with a higher occurrence of severe or gangrenous acute cholecystitis.

This explains the importance of stratifying geriatric patients as a means to select the proper surgical treatment. Evidence shows that cholecystectomy remains the main treatment for acute lithiasic cholecystitis, even in older patients,<sup>61</sup> and that mortality is significantly lower if surgery is performed during the same hospitalization.<sup>62</sup> Assessment of a patient's frailty is mandatory

alongside the related surgical risk, using validated scores, the mortality rates connected to surgical or conservative treatment, the probability of recurrence, and life expectancy. If a surgical approach is feasible, laparoscopic cholecystectomy appears to be the treatment of choice even in the elderly patient, as it is safe, a well-established procedure, it has low complication rates and reduced post-operative hospital stay.<sup>63</sup> The relatively higher rate of laparotomic conversions in older patients as reported in the literature,<sup>64,65</sup> is probably due to more severe cholecystitis upon admission.

If the patient is deemed "not fit" for surgery, the available alternatives are percutaneous cholecystostomy and antibiotic treatment.

Percutaneous cholecystostomy, either echo- or CT-guided, may be considered in patients over 65 years of age, with ASA scores III or IV, PF 3 or 4, or with septic shock who are considered unfit for surgery.<sup>13</sup> This treatment can be life-saving in emergency settings in those patients who would not tolerate immediate general anesthesia and surgery, but can also be considered as a bridge-to-surgery intervention.<sup>66,67,68</sup> The cholecystostomy catheter can be removed 4-6 weeks after its positioning, following a thorough study of the biliary duct with cholangiography.<sup>69,70</sup>

Empirical broad-spectrum antibiotic treatment may be reserved for those patients "not fit" for surgery, whose clinical, laboratory, or radiological findings are not striking enough to proceed with a percutaneous cholecystostomy.<sup>13</sup>

In the case of associated biliary lithiasis, the patient may undergo preoperative, intraoperative, or postoperative ERCP.<sup>71,72</sup>

### Acute appendicitis

Acute appendicitis is typical of the youth; only 10% of cases occur in patients aged 60 years or older,<sup>73</sup> in whom it is linked to more complications and mortality, due to frequent diagnostic and therapeutic delays. Divergent clinical presentation is recurrent in the geriatric patient and it includes atypical symptoms, late diagnosis, perforation, and septic complications.<sup>74,75</sup> In older patients, the rate of perforated appendicitis reaches up to 50-70% of cases, compared to 20-30% in the young.<sup>76,77</sup>

Typical clinical manifestations (fever, pain in the right abdominal quadrants, and leucocytosis) occur in only 10-26% of patients past 60 years of age;<sup>78</sup> this is due to age itself (reduced nerve sensitivity and abdominal musculature hyposensitivity), to the patient's comorbidities (*i.e.*, diabetes or cognitive decline) and the possible intake of pain-relieving or steroid drugs for other reasons, which increases the risk of perforated appendicitis upon admission.<sup>79</sup> Literature documents indeed that the perforation risk is much higher in patients over 80 years of age, even compared to patients aged between 60 and 79 years; this appears to be related to the common diagnostic and therapeutic delay linked to a reduced perception of pain, loneliness, comorbidities, limited physiological reserve, differential diagnosis with other typical geriatric diseases, and a higher chance of prior abdominal surgery: all these factors increase the overall surgical risk compared to younger patients. It has been observed that surgeons are often hesitant to operate on an 80-year-old patient and tend to favor non-operative treatment until explicit clinical-laboratory worsening: this delay only worsens the risk of complicated appendicitis.<sup>80</sup> Moreover, this setting requires a multidimensional evaluation of the geriatric patient to discern the best clinical and operative path taking advantage of the several scores that allow risk stratification, proper communication with the patient and family members, and receiving assistance from a geriatric consultant in post-operative period to decrease complications, length of stay and rate of institutionalization.

## Acute bowel obstruction

Key risk factors for the development of mechanical bowel obstruction are prior abdominal surgery (especially pelvic surgery), malignant pathology, hernias, volvulus, and recurrent diverticulitis; uncommon ones involve chronic inflammatory diseases and impacted stones. Diagnosis is based on clinical and radiological examinations, especially CT scans without and with i.v. contrast, which allows differential diagnoses.<sup>81,82</sup> In the literature, only a few scattered studies focus on geriatric emergency surgery due to intestinal obstruction, surgical strategies, and morbidity-mortality rates.<sup>83</sup> Early mortality rate (within 30 days after the surgical act) appears to be higher in older patients rather than the younger ones (9.1-23.5% in the elderly versus 0-3.2% in the young patient);<sup>83</sup> likewise, intra-hospital mortality was found to be around 18.2% in the geriatric patient and roughly 8.9% in the non-geriatric patient.<sup>84</sup> Length of stay is longer in older patients, with a greater risk of developing medical complications linked to hospitalization itself.<sup>85</sup> On the other hand, the resection rate, which is primarily related to vascular ischemia secondary to a protracted intestinal occlusion, has no statistically significant differences between the two groups.<sup>84</sup> Moreover, geriatric patients tend to be admitted to the Emergency Department at a worse and more frequently complicated stage of the pathology at issue, worsening de facto the surgical outcome compared to the non-geriatric patient, despite appropriate timing: clinical presentation delay decreases the efficiency of subsequent rehydrating treatment, electrolytes' management, and gastric decompression with a nasogastric tube, thus increasing the urgency of a surgical approach.<sup>84</sup> Furthermore, in geriatric patients, mechanical occlusion may have a more frequent neoplastic origin, with a higher rate of ostomy packing.<sup>84</sup> 30-day mortality is significantly higher in patients with mechanical occlusion of the large intestine (with evidence of intrinsic neoplastic origin in 60-70% of cases).<sup>81,86-90</sup>

## Mesenteric ischemia

Mesenteric ischemia can affect both the small and large bowel and is characterized by a mortality rate of 20-80% in the adult population regardless of age.<sup>91,92</sup> The main pathogenic factors include atherosclerosis, which is the most common cause in geriatric patients,<sup>93</sup> cardiac arrhythmias and valvular disease, vasoconstrictor drugs, and hereditary coagulation disorders.<sup>94,95</sup> The appearance of acute symptoms demands a speedy treatment, proceeding with an emergency laparotomy, possible revascularisation, and potential intestinal resection if needed.<sup>96</sup> According to the literature, 30-day mortality is around 31.5% in patients aged 65 years or older undergoing emergency surgery for mesenteric ischemia.<sup>97</sup> Usual complications include difficult weaning from the ventilator (30%), post-operative septic shock (22.4%), pneumonia (19.9%), and unexpected intubation (10.8%).<sup>97</sup> Preoperative factors that increase mortality and the risk of post-surgical complications are preoperative septic shock,<sup>92,97</sup> terminal renal failure undergoing dialysis,<sup>97</sup> reduced respiratory function<sup>97,98</sup> and a history of recent unintentional weight loss,<sup>97,99</sup> which can be considered as an indirect indicator of frailty and sarcopenia. It has been observed that without these risk factors mortality rate is drastically reduced, with a 30-day survival rate of 84% in patients from 65 to 80 years of age undergoing emergent surgical treatment for mesenteric ischemia.<sup>97</sup> Literature has demonstrated that mesenteric ischemia, even in geriatric patients, is no longer a sure death sentence, but requires individual risk stratification based on primary diagnosis, patient's age, specific preoperative comorbidities, and expected outcome.

## Treatment and palliation of surgical emergencies in geriatric patients with colorectal cancer

The incidence of colorectal cancer (CRC) increases with age: the average age of diagnosis is 68 years in men and 72 years in women.<sup>100</sup> More than one-third of cases are diagnosed in an emergency setting,<sup>101</sup> especially in older patients.<sup>102-104</sup> Advanced age, social deprivation, and belonging to an ethnic minority are the main risk factors for an emergency ED presentation of colorectal cancer.<sup>105</sup> Emergency treatment of CRC is generally associated with a worse prognosis, related to the recurrent presence of occlusion, perforation, and locally advanced or metastatic disease, leading thus increased morbidity and mortality.<sup>106</sup>

### Pre-operative assessment

The choice between palliative or radical curative treatment is crucial in the preoperative stage. The emergent nature of the situation often precludes a multidisciplinary assessment with an oncological discussion, so it is mandatory to acquire as much information as possible about the patient, his or her pathological history, lifestyle, social and family support, and associate it with clinical, laboratory, and radiological data, to estimate post-surgical morbidity and mortality, oncological prognosis and functional recovery in the medium and long term:<sup>40</sup> i) predicting post-surgical morbidity and mortality: in case of emergency admission for CRC, the major risk factors for adverse surgical outcomes are: proximal colon damage and/or peritonitis, being aged over 75 years old, comorbidities, advanced neoplastic stage, hemodynamic instability.<sup>107-111</sup> The prevailing risk factor is the patient's frailty.<sup>19</sup> The surgeon has many validated scores (e.g. P-POSSUM, fTRST, TEGS-FI, etc.) including the CT-scan assessed value of sarcopenia available to attain the most objective information as possible; ii) on the state of frailty and the consequent risk of postoperative morbidity and mortality;<sup>40</sup> iii) oncological prognosis: abdominal CT scan without and with contrast medium provides information on the patient's status (presence of perforation, peritonitis, sarcopenia, other pathologies), on the oncological stage (locoregional and distant neoplastic extension) and bowel status (dilation, possibility of endoscopic stent placement, choice of possible ostomy site).<sup>40</sup> The presence of perforated bowel is associated with a negative outcome and appears to expose the patient to an increased risk of peritoneal carcinosis (between 14% and 45% of cases);<sup>112</sup> iv) predicting functional recovery in the medium and long term and the consequences of a possible ostomy: functional recovery depends on the patient's preoperative morbidity and frailty, but also his/her social and family support. This is essential to understand what the patient's priorities and goals are, to determine what level of health status the patient would find acceptable, and to discuss palliative treatment and possible definitive ostomy placement with the patient and family members.<sup>113</sup>

### Surgical treatment

In the case of bowel perforation the primary goal is to control sepsis and only secondarily malignancy, performing a damage-control surgery in case of a hemodynamically unstable patient with a good performance status.<sup>40</sup>

In case of intestinal obstruction caused by colorectal cancer, the surgical treatment is affected by the neoplasm's site: a right colon localization can be treated surgically with right hemicolectomy and possible intestinal anastomosis if the neoplasm is resectable, or alternately with a decompressive ileostomy in locally advanced disease or patient instability;<sup>40</sup> a rectal location should be treated with

decompression with an ostomy or, if the neoplasm appears easily resectable, with a Hartmann sigma-rectal resection (stenting in rectal cancers is more commonly associated with pain and stent migration);<sup>40</sup> left colon localization may be treated with endoscopic stent placement or with ostomy as a bridge to surgery if a radical resection is programmed. The timing of the second surgical act may fluctuate from 7 to 10 days within the same hospitalization, or require a new hospitalization. In the case of palliative surgery, stent placement or ostomy may be definitive in the absence of any further surgical indications.<sup>114-118</sup>

For stage IV tumors, in patients with good PF, a resection of the primary tumor despite the absence of resectable metastases or a bridging treatment to demolitive resection may still be taken into consideration.<sup>119</sup> Evidence of peritoneal carcinosis diverts the surgeon's therapeutic choice towards a non-invasive treatment or a palliative ostomy.<sup>120</sup>

Both in the case of palliative and radical surgery, a laparoscopic approach should be preferred, as it is associated with lower pain and faster recovery in the postoperative period.<sup>121,122</sup>

### Post-operative management

Older patients admitted to the ED for colorectal cancer needing emergency surgery have a higher risk of post-operative morbidity and mortality than patients undergoing elective surgery.<sup>102,123,124</sup> Early mobilization, optimization of fluid therapy, early oral nutrition, and the use of non-opioid analgesics grant a faster patient recovery<sup>125,126</sup> and prevention of postoperative delirium,<sup>127</sup> in the meanwhile a personalized geriatric assessment after surgery can help the surgeon in the prevention and management of post-operative morbidity and in planning hospital discharge.<sup>128</sup>

### Damage control surgery

Damage-control surgery is widely recognized as a life-saving technique in unstable trauma patients or patients with severe surgical pathology treated in an emergency setting to allow further resuscitative maneuvers before undertaking definitive surgery.<sup>129</sup> The basic principles of damage-control surgery are the control of hemorrhage and peritoneal contamination, with a significant reduction in operating time, and the packing of a laparostomy, so that further resuscitative maneuvers can be performed in an intensive care setting and, once the patient is stabilized, proceed with curative surgery<sup>130</sup> 24-48 hours later.<sup>131</sup> The enactment of these damage-control surgery principles in older patients has demonstrated a reduction in mortality, ostomy rate, and time-lapse before subsequent ostomy closure.<sup>132-134</sup> Damage-control surgery followed by reintervention within 24 and 48 hours has not highlighted any significant difference in morbidity, mortality, or abdominal-wall closure rate in elderly patients compared to regular adult ones.<sup>135</sup> Damage-control surgery is thus feasible and should be considered in the geriatric patient after careful pre-operative assessment and patient stratification to avoid "unnecessary" surgical procedures in extremely frail and compromised patients for whom the line between necessary and inappropriate treatment may become very blurred.

### ERAS in emergency surgery in the elderly

The application of ERAS (Enhanced Recovery After Surgery) ideals in addition to a laparoscopic surgical approach in emergency surgery settings has shown benefits regarding morbidity, mortality, and length of stay in post-operative care.<sup>125</sup> In an emergency setting

the ERAS protocol is not fully applicable in its entirety, but some features, such as early mobilization, optimization of fluid therapy, early nutrition, and non-opioid analgesic therapy can be implemented in the post-operative setting to thwart complications and encourage rehabilitation.<sup>40</sup>

Early mobilization wards off loss of muscle mass and decreases pressure ulcers, deep vein thrombosis in the lower limbs, respiratory complications, and delirium, if supported by appropriate fall prevention protocols. Early post-operative rehabilitation programs have proven safe and effective in fending off and lowering functional decline in hospitalized acute geriatric patients, including those who have undertaken emergent abdominal surgery.<sup>136,137</sup>

Optimization of fluid therapy, along with a daily input/output fluid balance is crucial, especially in patients undergoing massive intraoperative liquid infusions. Geriatric patients encounter on one side a high risk of hypervolemia and congestive heart failure, paralytic ileus, and edema;<sup>46</sup> on the other side, dehydration is associated with acute renal failure and delirium.<sup>138</sup>

Nutrition plays a cardinal role in the geriatric patient's rehabilitation. These patients often present with malnutrition (5.8-50.5%)<sup>139</sup> due to long periods of fasting or inappetence. The combination of metabolic disorders, malnutrition, and sarcopenia relates to a limited ability to respond properly to surgical stress in the short, medium, and long term.<sup>140,141</sup> One of the key goals is thus to restore oral food intake as soon as possible and to potentially start supportive nutritional therapy should any surgical risks arise.<sup>141</sup>

### Conclusions

The physical, psychological, and social characteristics of a geriatric patient depict him or her as more fragile and vulnerable. Advanced age is associated with an increased risk of comorbidity and loss of autonomy, as well as a decrease in life expectancy. Geriatric age does not necessarily correlate with frailty, which indeed has a linear, but not univocal, trend with age. Frailty is also associated with an increased risk of all complications and death, after both elective surgery and emergency surgery. Yet, despite the ability of elective surgery to assess the geriatric patient in a multidisciplinary manner and properly prepare him/her for the upcoming surgical stress, this is not feasible in acute situations, where the time factor is crucial. Therefore, several fragility scores have been developed over the years to allow a preoperative assessment of the urgent surgical patient, to estimate the risk of morbidity and mortality, and to guide the surgeon toward the proper therapeutic or palliative treatment and limit the risk of complications in the postoperative period through the adoption of measures tailored to each patient. Many studies have managed to appraise the benefits of the different scores in an emergency setting in the geriatric patient, whereas there are still only a handful of studies to have analyzed the outcomes of a specific urgent surgical procedure performed in the geriatric patient applying one or more specific scores. These considerations certainly give rise to further pondering and discussion, given the constant aging of the general population and the increase in life expectancy even in frail patients.

### References

1. Kapadia M, Obaid O, Nelson A, et al. Evaluation of frailty assessment compliance in acute care surgery: changing trends, lessons learned. *J Surg Res* 2022;270:236-44.
2. Panayi AC, Orkaby AR, Sakthivel D, et al. Impact of frailty on

- outcomes in surgical patients: A systematic review and meta-analysis. *Am J Surg* 2019;218:393-400.
3. Trotter G. Mediating disputes about medical futility. *Camb Q Health Ethics* 1999;8:527-37.
  4. Javanmard-Emamghissi H, Moug SJ. The virtual uncertainty of futility in emergency surgery. *Br J Surg* 2022;109:1184-5.
  5. Adelson BH. Resolving ethical dilemmas: a guide for clinicians. *JAMA* 1995;274:587-88.
  6. Parmar KL, Law J, Carter B, et al. Frailty in older patients undergoing emergency laparotomy: results from the UK observational Emergency Laparotomy and Frailty (ELF) study. *Ann Surg* 2021;273:709-18.
  7. Law J, Welch C, Javanmard-Emamghissi H, et al. Decision-making for older patients undergoing emergency laparotomy: defining patient and clinician values and priorities. *Colorectal Dis* 2020;22:1694-703.
  8. Lambden JP, Chamberlin P, Kozlov E, et al. Association of perceived futile or potentially inappropriate care with burnout and thoughts of quitting among health-care providers. *Am J Hosp Palliat Care* 2019;36:200-6.
  9. Aggarwal G, Scott M, Peden CJ. Emergency laparotomy. *Anesthesiol Clin* 2022;40:199-211.
  10. Ablett AD, McCarthy K, Carter B. A practical risk scale for predicting morbidity and mortality in the emergency general surgical setting: A prospective multi-center study. *Int J Surg* 2018;60:236-44.
  11. Ho VP, Bensken WP, Warner DF. Association of complex multimorbidity and long-term survival after emergency general surgery in older patients with medicare. *JAMA Surg* 2022;157:499-506.
  12. Aucoin S, McIsaac DI. Emergency general surgery in older adults: a review. *Anesthesiol Clin* 2019;37:493-505.
  13. Pisano M, Ceresoli M, Cimbanassi S, et al. 2017 WSES and SICG guidelines on acute calculous cholecystitis in elderly population. *World J Emerg Surg* 2019;14:10.
  14. Elamin A, Tsoutsanis P, Sinan L. Emergency general surgery: predicting morbidity and mortality in the geriatric population. *Surg J (NY)* 2022;8:e270-8.
  15. Ahmed SM, Alali MA, Atuel KLG. American Society of Anesthesiologists Physical Status Classification System: history, development, reliability, and its future. In: Sozen S, editor. *Surgical Recovery*. London: IntechOpen; 2019. Available from: <https://www.intechopen.com/chapters/70432> doi: 10.5772/intechopen.89704
  16. Bolger JC, Zaidi A, Fuentes-Bonachera A, Kelly ME. Emergency surgery in octogenarians: Outcomes and factors affecting mortality in the general hospital setting. *Geriatr Gerontol Int* 2018;18:1211-4.
  17. Vilches-Moraga A, Rowley M, Fox J. Emergency laparotomy in the older patient: factors predictive of 12-month mortality-Salford-POPS-GS. An observational study. *Aging Clin Exp Res* 2020;32:2367-73.
  18. Sezen Ö, Çevik B. Emergency surgery in geriatrics: A retrospective evaluation in a single center. *Ulus Travma Acil Cerrahi Derg* 2020;26:411-7.
  19. Hewitt J, Carter B, McCarthy K. Frailty predicts mortality in all emergency surgical admissions regardless of age. An observational study. *Age Ageing* 2019;48:388-94.
  20. Zattoni D, Montroni I, Saur NM, et al. Prediction of functional loss in emergency surgery is possible with a simple frailty screening tool. *World J Emerg Surg* 2021;16:12.
  21. Fehlmann CA, Patel D, McCallum J. Association between mortality and frailty in emergency general surgery: a systematic review and meta-analysis. *Eur J Trauma Emerg Surg* 2022;48:141-51.
  22. Rockwood K, Song X, MacKnight C, et al. A global clinical measure of fitness and frailty in elderly people. *CMAJ* 2005;173:489-95.
  23. Goeteyn J, Evans LA, De Cleyn S, et al. Frailty as a predictor of mortality in the elderly emergency general surgery patient. *Acta Chir Belg* 2017;117:370-5.
  24. Eamer GJ, Clement F, Holroyd-Leduc J, et al. Frailty predicts increased costs in emergent general surgery patients: A prospective cohort cost analysis. *Surgery* 2019;166:82-7.
  25. Cheung A, Haas B, Ringer TJ, et al. Canadian study of health and aging clinical frailty scale: does it predict adverse outcomes among geriatric trauma patients? *J Am Coll Surg* 2017;225:658-665.e3.
  26. Mintziras I, Miligkos M, Wächter S, et al. Sarcopenia and sarcopenic obesity are significantly associated with poorer overall survival in patients with pancreatic cancer: Systematic review and meta-analysis. *Int J Surg* 2018;59:19-26.
  27. Kołodziejka K, Witowski J, Tylec P, et al. Radiological features for frailty assessment in patients requiring emergency laparotomy. *J Clin Med* 2022;11:5365.
  28. Thu K, Nguyen HPT, Gogulan T, et al. Care of older people in surgery for general surgery: a single centre experience. *ANZ J Surg* 2021;91:890-5.
  29. Kaplan SJ, Pham TN, Arbabi S, et al. Association of radiologic indicators of frailty with 1-year mortality in older trauma patients: opportunistic screening for sarcopenia and osteopenia. *JAMA Surg* 2017;152:e164604.
  30. Pędziwiatr M, Pisarska M, Major P, et al. Laparoscopic colorectal cancer surgery combined with enhanced recovery after surgery protocol (ERAS) reduces the negative impact of sarcopenia on short-term outcomes. *Eur J Surg Oncol* 2016;42:779-87.
  31. Pecorelli N, Capretti G, Sandini M, et al. Impact of sarcopenic obesity on failure to rescue from major complications following pancreaticoduodenectomy for cancer: results from a multicenter study. *Ann Surg Oncol* 2018;25:308-17.
  32. Ratnayake CBB, Wells C, Olsson M, et al. Sarcopenic obesity and post-operative morbidity after pancreatic surgery: a cohort study. *ANZ J Surg* 2019;89:1587-92.
  33. Zhang WT, Lin J, Chen WS, et al. Sarcopenic obesity is associated with severe postoperative complications in gastric cancer patients undergoing gastrectomy: a prospective study. *J Gastrointest Surg* 2018;22:1861-9.
  34. Mason MC, Crees AL, Dean MR, Bashir N. Establishing a proactive geriatrician led comprehensive geriatric assessment in older emergency surgery patients: Outcomes of a pilot study. *Int J Clin Pract* 2018;72:e13096.
  35. Khan M, Jehan F, Zeeshan M, et al. Failure to rescue after emergency general surgery in geriatric patients: does frailty matter? *J Surg Res* 2019;233:397-402.
  36. Palaniappan S, Soiza RL, Duffy S, et al. Comparison of the clinical frailty score (CFS) to the National Emergency Laparotomy Audit (NELA) risk calculator in all patients undergoing emergency laparotomy. *Colorectal Dis* 2022;24:782-9.
  37. Aitken RM, Partridge JSL, Oliver CM, et al. Older patients undergoing emergency laparotomy: observations from the National Emergency Laparotomy Audit (NELA) years 1-4. *Age Ageing* 2020;49:656-63.
  38. Ho VP, Schiltz NK, Reimer AP, et al. High-risk comorbidity combinations in older patients undergoing emergency general surgery. *J Am Geriatr Soc* 2019;67:503-10.
  39. Torrance AD, Powell SL, Griffiths EA. Emergency surgery in



- the elderly: challenges and solutions. *Open Access Emerg Med* 2015;7:55-68.
40. Zattoni D, Christoforidis D. How best to palliate and treat emergency conditions in geriatric patients with colorectal cancer. *Eur J Surg Oncol* 2020;46:369-78.
  41. Tan HL, Chia STX, Nadkarni NV, et al. Frailty and functional decline after emergency abdominal surgery in the elderly: a prospective cohort study. *World J Emerg Surg* 2019;14:62.
  42. Murphy PB, Savage SA, Zarzaur BL. Impact of patient frailty on morbidity and mortality after common emergency general surgery operations. *J Surg Res* 2020;247:95-102.
  43. Becher RD, Murphy TE, Gahbauer EA, et al. Factors associated with functional recovery among older survivors of major surgery. *Ann Surg* 2020;272:92-8.
  44. Liu X, Le MK, Lim AYC, et al. Perspectives on frailty screening, management and its implementation among acute care providers in Singapore: a qualitative study. *BMC Geriatr* 2022;22:58.
  45. Kenawy DM, Renshaw SM, George E, et al. Increasing frailty in geriatric emergency general surgery: a cause for concern. *J Surg Res* 2021;266:320-7.
  46. Zattoni D, Montroni I, Saur NM, et al. A simple screening tool to predict outcomes in older adults undergoing emergency general surgery. *J Am Geriatr Soc* 2019;67:309-16.
  47. Engelhardt KE, Reuter Q, Liu J, et al. Frailty screening and a frailty pathway decrease length of stay, loss of independence, and 30-day readmission rates in frail geriatric trauma and emergency general surgery patients. *J Trauma Acute Care Surg* 2018;85:167-73.
  48. Mahmooth Z, Carpenter E, Lin E, et al. Frailty assessment in the acute care surgery population - the agreement and predictive value on length of stay and re-admission of 3 different instruments in a prospective cohort. *Am J Surg* 2020;220:1058-63.
  49. Orouji Jokar T, Ibraheem K, Rhee P, et al. Emergency general surgery specific frailty index: A validation study. *J Trauma Acute Care Surg* 2016;81:254-60.
  50. Weiss HK, Stocker BW, Weingarten N, et al. Electronic medical record versus bedside assessment: how to evaluate frailty in trauma and emergency general surgery patients? *J Surg Res* 2020;246:464-75.
  51. Boyd-Carson H, Gana T, Lockwood S, et al. A review of surgical and peri-operative factors to consider in emergency laparotomy care. *Anaesthesia* 2020;75:e75-e82.
  52. Ablett AD, McCarthy K, Carter B, et al. A practical risk scale for predicting morbidity and mortality in the emergency general surgical setting: A prospective multi-center study. *Int J Surg* 2018;60:236-44.
  53. Festi D, Dormi A, Capodicasa S, et al. Incidence of gallstone disease in Italy: results from a multicenter, population-based Italian study (the MICOL project). *World J Gastroenterol* 2008;14:5282-9.
  54. Sanson TG, O'Keefe KP. Evaluation of abdominal pain in the elderly. *Emerg Med Clin North Am* 1996;14:615-27.
  55. Ragsdale L, Southerland L. Acute abdominal pain in the older adult. *Emerg Med Clin North Am* 2011;29:429-48, x.
  56. Ross SO, Forsmark CE. Pancreatic and biliary disorders in the elderly. *Gastroenterol Clin North Am* 2001;30:531-45, x.
  57. Lavretsky H, Newhouse PA. Stress, inflammation, and aging. *Am J Geriatr Psychiatry* 2012;20:729-33.
  58. Parker LJ, Vukov LF, Wollan PC. Emergency department evaluation of geriatric patients with acute cholecystitis. *Acad Emerg Med* 1997;4:51-5.
  59. McGillicuddy EA, Schuster KM, Brown E, et al. Acute cholecystitis in the elderly: use of computed tomography and correlation with ultrasonography. *Am J Surg* 2011;202:524-7.
  60. Asiltürk Lülleci Z, Başyığıt S, Pirinççi Salmaz F, et al. Comparison of ultrasonographic and laboratory findings of acute cholecystitis between elderly and nonelderly patients. *Turk J Med Sci* 2016;46:1428-33.
  61. Preston SD, Southall AR, Nel M, Das SK. Geriatric surgery is about disease, not age. *J R Soc Med* 2008;101:409-15.
  62. Riall TS, Zhang D, Townsend CM Jr, et al. Failure to perform cholecystectomy for acute cholecystitis in elderly patients is associated with increased morbidity, mortality, and cost. *J Am Coll Surg* 2010;210:668-77, 677-9.
  63. Coccolini F, Catena F, Pisano M, et al. Open versus laparoscopic cholecystectomy in acute cholecystitis. Systematic review and meta-analysis. *Int J Surg* 2015;18:196-204. Erratum in: *Int J Surg* 2015;24:107.
  64. Elshaer M, Gravante G, Thomas K, et al. Subtotal cholecystectomy for "difficult gallbladders": systematic review and meta-analysis. *JAMA Surg* 2015;150:159-68.
  65. Palanivelu C, Rajan PS, Jani K, et al. Laparoscopic cholecystectomy in cirrhotic patients: the role of subtotal cholecystectomy and its variants. *J Am Coll Surg* 2006;203:145-51.
  66. Ansaloni L, Pisano M, Coccolini F, et al. 2016 WSES guidelines on acute calculous cholecystitis. *World J Emerg Surg* 2016;11:25.
  67. Bakkaloglu H, Yanar H, Guloglu R, et al. Ultrasound guided percutaneous cholecystostomy in high-risk patients for surgical intervention. *World J Gastroenterol* 2006;12:7179-82.
  68. Miura F, Takada T, Kawarada Y, et al. Flowcharts for the diagnosis and treatment of acute cholangitis and cholecystitis: Tokyo Guidelines. *J Hepatobiliary Pancreat Surg* 2007;14:27-34.
  69. Nasim S, Khan S, Alvi R, Chaudhary M. Emerging indications for percutaneous cholecystostomy for the management of acute cholecystitis--a retrospective review. *Int J Surg* 2011;9:456-9.
  70. Venara A, Carretier V, Lebigot J, Lermite E. Technique and indications of percutaneous cholecystostomy in the management of cholecystitis in 2014. *J Visc Surg* 2014;151:435-9.
  71. Dasari BV, Tan CJ, Gurusamy KS, et al. Surgical versus endoscopic treatment of bile duct stones. *Cochrane Database Syst Rev* 2013;2013:CD003327.
  72. Wang B, Guo Z, Liu Z, et al. Preoperative versus intraoperative endoscopic sphincterotomy in patients with gallbladder and suspected common bile duct stones: system review and meta-analysis. *Surg Endosc* 2013;27:2454-65.
  73. Blomqvist PG, Andersson RE, Granath F, et al. Mortality after appendectomy in Sweden, 1987-1996. *Ann Surg* 2001;233:455-60.
  74. Massarweh NN, Legner VJ, Symons RG, et al. Impact of advancing age on abdominal surgical outcomes. *Arch Surg* 2009;144:1108-14.
  75. Lee JF, Leow CK, Lau WY. Appendicitis in the elderly. *Aust N Z J Surg* 2000;70:593-6.
  76. Yamini D, Vargas H, Bongard F, et al. Perforated appendicitis: is it truly a surgical urgency? *Am Surg* 1998;64:970-5.
  77. Sanson TG, O'Keefe KP. Evaluation of abdominal pain in the elderly. *Emerg Med Clin North Am* 1996;14:615-27.
  78. Paranjape C, Dalia S, Pan J, Horattas M. Appendicitis in the elderly: a change in the laparoscopic era. *Surg Endosc* 2007;21:777-81.
  79. Omari AH, Khammash MR, Qasaimeh GR, et al. Acute appendicitis in the elderly: risk factors for perforation. *World J Emerg Surg* 2014;9:6.

80. Tan KY, Chen CM, Ng C, et al. Which octogenarians do poorly after major open abdominal surgery in our Asian population? *World J Surg* 2006;30:547-52.
81. Ten Broek RPG, Krielen P, Di Saverio S, et al. Bologna guidelines for diagnosis and management of adhesive small bowel obstruction (ASBO): 2017 update of the evidence-based guidelines from the world society of emergency surgery ASBO working group. *World J Emerg Surg* 2018;13:24.
82. Branco BC, Bamparas G, Schnüriger B, et al. Systematic review and meta-analysis of the diagnostic and therapeutic role of water-soluble contrast agent in adhesive small bowel obstruction. *Br J Surg* 2010;97:470-8.
83. Krause WR, Webb TP. Geriatric small bowel obstruction: an analysis of treatment and outcomes compared with a younger cohort. *Am J Surg* 2015;209:347-51.
84. Paul CJJ, Dohmen J, van Beekum CJ, et al. Surgical treatment of mechanical bowel obstruction: characteristics and outcomes of geriatric patients compared to a younger cohort. *Int J Colorectal Dis* 2022;37:1281-8.
85. van Beekum CJ, Stoffels B, von Websky M, et al. Der mechanische Ileus bei geriatrischen Patienten: Ätiologie und perioperative Morbidität/Mortalität im Vergleich zu einer jüngeren Patientenkohorte Mechanical bowel obstruction in geriatric patients : Etiology and perioperative morbidity/mortality compared with a younger cohort. *Med Klin Intensivmed Notfmed* 2020;115:22-28. German.
86. Cappell MS, Batke M. Mechanical obstruction of the small bowel and colon. *Med Clin North Am* 2008;92:575-97.
87. Karakaş DÖ, Yeşiltaş M, Gökçek B, et al. Etiology, management, and survival of acute mechanical bowel obstruction: Five-year results of a training and research hospital in Turkey. *Ulus Travma Acil Cerrahi Derg* 2019;25:268-80.
88. Santangelo ML, Grifasi C, Criscitiello C, et al. Bowel obstruction and peritoneal carcinomatosis in the elderly. A systematic review. *Aging Clin Exp Res* 2017;29:73-8.
89. Bükki J. Re: recommendations for bowel obstruction with peritoneal carcinomatosis by laval et Al. *J Pain Symptom Manage* 2014;48:e5-6.
90. Krstić S, Resanovic V, Alempijević T, et al. Hartmann's procedure vs loop colostomy in the treatment of obstructive rectosigmoid cancer. *World J Emerg Surg* 2014;9:52.
91. Crawford RS, Harris DG, Klyushnenkova EN, et al. A statewide analysis of the incidence and outcomes of acute mesenteric ischemia in Maryland from 2009 to 2013. *Front Surg* 2016;3:22.
92. Tseng J, Loper B, Jain M, et al. Predictive factors of mortality after colectomy in ischemic colitis: an ACS-NSQIP database study. *Trauma Surg Acute Care Open* 2017;2:e000126.
93. Roobottom CA, Dubbins PA. Significant disease of the celiac and superior mesenteric arteries in asymptomatic patients: predictive value of Doppler sonography. *AJR Am J Roentgenol* 1993;161:985-8.
94. Bekwelem W, Connolly SJ, Halperin JL, et al. Extracranial systemic embolic events in patients with nonvalvular atrial fibrillation: incidence, risk factors, and outcomes. *Circulation* 2015;132:796-803.
95. Amitrano L, Brancaccio V, Guardascione MA, et al. High prevalence of thrombophilic genotypes in patients with acute mesenteric vein thrombosis. *Am J Gastroenterol* 2001;96:146-9.
96. Bala M, Kashuk J, Moore EE, et al. Acute mesenteric ischemia: guidelines of the World Society of Emergency Surgery. *World J Emerg Surg* 2017;12:38.
97. Gebran A, Gallastegi AD, Fagenholz P, et al. Factors that predict survival from mesenteric ischemia in the geriatric patient: not a death sentence. *Am Surg* 2021;31348211050585.
98. Kleinhenz ME, Lewis CY. Chronic ventilator dependence in elderly patients. *Clin Geriatr Med* 2000;16:735-56.
99. Bales CW, Ritchie CS. Sarcopenia, weight loss, and nutritional frailty in the elderly. *Annu Rev Nutr* 2002;22:309-23.
100. Viale PH. The American Cancer Society's Facts & Figures: 2020 Edition. *J Adv Pract Oncol* 2020;11:135-6.
101. Baer C, Menon R, Bastawrous S, Bastawrous A. Emergency presentations of colorectal cancer. *Surg Clin North Am* 2017;97:529-45.
102. Neuman HB, O'Connor ES, Weiss J, et al. Surgical treatment of colon cancer in patients aged 80 years and older: analysis of 31,574 patients in the SEER-Medicare database. *Cancer* 2013;119:639-47.
103. Esteva M, Ruidiáz M, Sánchez MA, et al. Emergency presentation of colorectal patients in Spain. *PLoS One* 2018;13:e0203556. Erratum in: *PLoS One* 2018;13:e0207143.
104. National Bowel Cancer Audit, Annual report. 2018. Available from: [https://www.hqip.org.uk/wp-content/uploads/2018/12/ref73\\_nboca-annual-report-2018\\_final\\_20181213.pdf](https://www.hqip.org.uk/wp-content/uploads/2018/12/ref73_nboca-annual-report-2018_final_20181213.pdf); 2018 Accessed 02/02/2019
105. Askari A, Nachiappan S, Currie A, et al. Who requires emergency surgery for colorectal cancer and can national screening programmes reduce this need? *Int J Surg* 2017;42:60-8.
106. Amri R, Bordeianou LG, Sylla P, Berger DL. Colon cancer surgery following emergency presentation: effects on admission and stage-adjusted outcomes. *Am J Surg* 2015;209:246-53.
107. Tekkis PP, Kinsman R, Thompson MR, et al. The Association of Coloproctology of Great Britain and Ireland study of large bowel obstruction caused by colorectal cancer. *Ann Surg* 2004;240:76-81.
108. Aslar AK, Ozdemir S, Mahmoudi H, Kuzu MA. Analysis of 230 cases of emergent surgery for obstructing colon cancer--lessons learned. *J Gastrointest Surg* 2011;15:110-9.
109. Alvarez JA, Baldonado RF, Bear IG, et al. Obstructing colorectal carcinoma: outcome and risk factors for morbidity and mortality. *Dig Surg* 2005;2:174-81.
110. Biondo S, Parés D, Frago R, et al. Large bowel obstruction: predictive factors for postoperative mortality. *Dis Colon Rectum* 2004;47:1889-97.
111. Manceau G, Mege D, Bridoux V, et al. Thirty-day mortality after emergency surgery for obstructing colon cancer: survey and dedicated score from the French Surgical Association. *Colorectal Dis* 2019;21:782-90.
112. Honoré C, Goéré D, Souadka A, et al. Definition of patients presenting a high risk of developing peritoneal carcinomatosis after curative surgery for colorectal cancer: a systematic review. *Ann Surg Oncol* 2013;20:183-92.
113. Cooper Z, Courtwright A, Karlage A, et al. Pitfalls in communication that lead to nonbeneficial emergency surgery in elderly patients with serious illness: description of the problem and elements of a solution. *Ann Surg* 2014;260:949-57.
114. Sebastian S, Johnston S, Geoghegan T, et al. Pooled analysis of the efficacy and safety of self-expanding metal stenting in malignant colorectal obstruction. *Am J Gastroenterol* 2004;99:2051-7.
115. Foo CC, Poon SHT, Chiu RHY, et al. Is bridge to surgery stenting a safe alternative to emergency surgery in malignant colonic obstruction: a meta-analysis of randomized control trials. *Surg Endosc* 2019;33:293-302.
116. Sabbagh C, Browet F, Diouf M, et al. Is stenting as "a bridge to surgery" an oncologically safe strategy for the management

- of acute, left-sided, malignant, colonic obstruction? A comparative study with a propensity score analysis. *Ann Surg* 2013;258:107-15.
117. van Hooft JE, Veld JV, Arnold D, et al. Self-expandable metal stents for obstructing colonic and extracolonic cancer: European Society of Gastrointestinal Endoscopy (ESGE) Guideline - Update 2020. *Endoscopy* 2020;52:389-407.
  118. Ansaloni L, Andersson RE, Bazzoli F, et al. Guidelienines in the management of obstructing cancer of the left colon: consensus conference of the world society of emergency surgery (WSES) and peritoneum and surgery (PnS) society. *World J Emerg Surg* 2010;5:29.
  119. Tarantino I, Warschkow R, Worni M, et al. Prognostic relevance of palliative primary tumor removal in 37,793 metastatic colorectal cancer patients: a population-based, propensity score-adjusted trend analysis. *Ann Surg* 2015;262:112-20.
  120. de Boer NL, Hagemans JAW, Schultze BTA, et al. Acute malignant obstruction in patients with peritoneal carcinomatosis: The role of palliative surgery. *Eur J Surg Oncol* 2019;45:389-93.
  121. Frasson M, Braga M, Vignali A, Zuliani W, Di Carlo V. Benefits of laparoscopic colorectal resection are more pronounced in elderly patients. *Dis Colon Rectum* 2008;51:296-300.
  122. Cerdán Santacruz C, Frasson M, Flor-Lorente B, et al. Laparoscopy may decrease morbidity and length of stay after elective colon cancer resection, especially in frail patients: results from an observational real-life study. *Surg Endosc* 2017;31:5032-42.
  123. Desserud KF, Veen T, Søreide K. Emergency general surgery in the geriatric patient. *Br J Surg* 2016;103:e52-61.
  124. Duron JJ, Duron E, Dugue T, et al. Risk factors for mortality in major digestive surgery in the elderly: a multicenter prospective study. *Ann Surg* 2011;254:375-82.
  125. Spanjersberg WR, van Sambeek JD, Bremers A, et al. Systematic review and meta-analysis for laparoscopic versus open colon surgery with or without an ERAS programme. *Surg Endosc* 2015;29:3443-53.
  126. Lohsiriwat V, Jitmongngan R. Enhanced recovery after surgery in emergency colorectal surgery: Review of literature and current practices. *World J Gastrointest Surg* 2019;11:41-52.
  127. Oh ES, Fong TG, Hshieh TT, Inouye SK. Delirium in older persons: advances in diagnosis and treatment. *JAMA* 2017;318:1161-74.
  128. Berian JR, Mohanty S, Ko CY, et al. Association of loss of independence with readmission and death after discharge in older patients after surgical procedures. *JAMA Surg* 2016;151:e161689.
  129. Holcomb JB, Jenkins D. Damage control resuscitation: directly addressing the early coagulopathy of trauma. *J Trauma* 2007;62:307-10.
  130. Rotondo MF, Schwab CW, McGonigal MD, et al. 'Damage control': an approach for improved survival in exsanguinating penetrating abdominal injury. *J Trauma* 1993;35:375-82; discussion 382-3.
  131. Sohn M, Agha A, Heitland W, et al. Damage control strategy for the treatment of perforated diverticulitis with generalized peritonitis. *Tech Coloproctol* 2016;20:577-83.
  132. Stawicki SP, Brooks A, Bilski T, et al. The concept of damage control: extending the paradigm to emergency general surgery. *Injury* 2008;39:93-101.
  133. Tartaglia D, Costa G, Camillò A, et al. Damage control surgery for perforated diverticulitis with diffuse peritonitis: saves lives and reduces ostomy. *World J Emerg Surg* 2019;14:19.
  134. Kafka-Ritsch R, Birkfellner F, Perathoner A, et al. Damage control surgery with abdominal vacuum and delayed bowel reconstruction in patients with perforated diverticulitis Hinchey III/IV. *J Gastrointest Surg* 2012;16:1915-22.
  135. Iacono SA, Krumrei NJ, Niroomand A, et al. Age is but a number: damage control surgery outcomes in geriatric emergency general surgery. *J Surg Res* 2021;267:452-7.
  136. McComb A, Warkentin LM, McNeely ML, Khadaroo RG. Development of a reconditioning program for elderly abdominal surgery patients: the Elder-friendly Approaches to the Surgical Environment-BEside reconditioning for Functional ImprovementTs (EASE-BE FIT) pilot study. *World J Emerg Surg* 2018;13:21.
  137. Martínez-Velilla N, Casas-Herrero A, Zambom-Ferraresi F, et al. Effect of exercise intervention on functional decline in very elderly patients during acute hospitalization: a randomized clinical trial. *JAMA Intern Med* 2019;179:28-36. Erratum in: *JAMA Intern Med* 2019;179:127.
  138. El-Sharkawy AM, Sahota O, Maughan RJ, Lobo DN. The pathophysiology of fluid and electrolyte balance in the older adult surgical patient. *Clin Nutr* 2014;33:6-13.
  139. Kaiser MJ, Bauer JM, Rämisch C. Frequency of malnutrition in older adults: a multinational perspective using the mini nutritional assessment. *J Am Geriatr Soc* 2010;58:1734-8.
  140. Rangel EL, Rios-Diaz AJ, Uyeda JW, et al. Sarcopenia increases risk of long-term mortality in elderly patients undergoing emergency abdominal surgery. *J Trauma Acute Care Surg* 2017;83:1179-86.
  141. Weimann A, Braga M, Carli F, et al. ESPEN guideline: Clinical nutrition in surgery. *Clin Nutr* 2017;36:623-50.